

Post-Operative Drop Instructions

There will be a total of *three* different bottles of drops to use after surgery. Two of these drops are the same ones you used before surgery and one is new. There are different brands that may be used. Use the charts below to locate the specific drops prescribed by your surgeon.

Note: You may use the drops together but wait 5 minutes between drops

Steroid Eye Drop Instructions:

	Week #1	Week #2	Week #3	Week #4
Prednisolone 1%	4 times a day	3 times a day	2 times a day	1 time a day
Lotemax	4 times a day	3 times a day	2 times a day	1 time a day
Durezol	4 times a day	2 times a day	2 times a day	1 time a day

Non-steroidal Eye Drop Instructions:

	Week #1	Week #2	Week #3	Week #4
Bromday	1 time a day	1 time a day	1 time a day	1 time a day
Xibrom	2 times a day	2 times a day	2 times a day	2 times a day
Acuvail	2 times a day	2 times a day	2 times a day	2 times a day
Nevanac	3 times a day	3 times a day	3 times a day	3 times a day
Ketorolac	4 times a day	4 times a day	4 times a day	4 times a day

Antibiotic Eye Drop Instructions:

	Week #1	Week #2	Week #3	Week #4
Zymaxid	2 times a day	NONE	NONE	NONE
Besivance	2 times a day	NONE	NONE	NONE
Vigamox	3 times a day	NONE	NONE	NONE
Ciprofloxacin	4 times a day	NONE	NONE	NONE